

THE X-RAY LUMINOSITY FUNCTION OF NEARBY RICH AND POOR CLUSTERS OF GALAXIES: A COSMOLOGICAL PROBE

MICHAEL J. LEDLOW¹, CHRIS LOKEN², JACK O. BURNS², FRAZER N. OWEN³,
AND WOLFGANG VOGES⁴

¹Institute for Astrophysics, University of New Mexico, Albuquerque, NM 87131

²Office of Research and Dept. of Physics & Astronomy, University of Missouri
Columbia, MO 65211

³National Radio Astronomy Observatory¹, Socorro, NM 87801

⁴Max Planck-Institut für Extraterrestrische Physik, Postfach 1603, D-85740,
Garching bei München, Germany

Accepted for publication in ApJ Letters

ABSTRACT

In this letter, we present a new determination of the local ($z \leq 0.09$) X-ray luminosity function (XLF) using a large, statistical sample of 294 Abell clusters and the *ROSAT* All-Sky-Survey. Despite the optical selection of this catalog, we find excellent agreement with other recent determinations of the local XLF. Given our large sample size, we have reduced errors by $\approx$ a factor of two for $L_{X(0.5-2.0 \text{ keV})} \geq 10^{43} h_{50}^{-2} \text{ ergs/sec}$. We combine our data with previous work to produce the most tightly constrained local determination of the XLF over three orders of magnitude in L_X in order to explore possible constraints imposed by the shape of the XLF on cosmological models. A set of currently viable cosmologies is used to construct theoretical XLFs assuming $L \propto M^p$ and a $\sigma_8 - \Omega_0$ constraint (from Viana & Liddle 1996) based on the local X-ray temperature function. We fit these models to our observed XLF and verify that the simplest adiabatic, analytic scaling relation (*e.g.* Kaiser 1986) disagrees strongly with observations. If we assume that clusters can be described by the pre-heated, constant core entropy models of Evrard & Henry (1991) then the observed XLF is consistent only with $0.1 < \Omega_0 < 0.4$ if the energy per unit mass in galaxies is roughly equal to the gas energy (*i.e.*, if $\beta \sim 1$).

Subject headings: cosmology: observations – galaxies: clusters: general – X-rays: general

1. INTRODUCTION AND BACKGROUND

Much of the work on the luminosity distributions of rich clusters has been motivated by the results of Henry *et al.* (1992) who found evidence for statistically significant negative evolution in the XLF (*i.e.* fewer high L_X clusters at higher z) at $z \geq 0.3$ for $L_{X(0.3-3.5 \text{ keV})} \geq 5 \times 10^{44} h_{50}^{-2} \text{ ergs sec}^{-1}$ from 67 clusters in the *Einstein* Extended Medium-Sensitivity Survey (EMSS). Recently, Vikhlinin *et al.* (1998) have confirmed the EMSS result at $z > 0.3$ for $L_{X(0.5-2 \text{ keV})} > 3 \times 10^{44} h_{50}^{-2} \text{ ergs/sec}$ from a 160 deg^2 survey from pointed *ROSAT* fields. They found a factor of 3-4 decrease in the number density of these high L_X clusters as compared to a zero-evolution model. Several other studies have claimed no evolution in the XLF out to redshifts as high as $z = 0.8$ (Burke *et al.* 1997; Jones *et al.* 1998; Rosati *et al.* 1998). However, none of these studies have sufficiently large search volumes to address evolution in the XLF at the highest X-ray luminosities and thus do not contradict the original EMSS result.

Of prime importance in any evolutionary study is an accurate determination of the *local* XLF as a baseline to compare with the distant cluster XLF. Until recently, even the local XLF was quite poorly constrained due to low cluster numbers. The largest local samples compiled to date are the X-ray Brightest Abell Clusters (XBACS) (Ebeling *et al.* 1993, 1996) and the Brightest Cluster Sample (BCS) of Ebeling *et al.* (1997, 1998). The BCS includes 199 X-

ray selected clusters down to $\approx 5 \times 10^{42} \text{ ergs sec}^{-1}$ in the 0.1 – 2.4 keV band out to $z \leq 0.3$. Consistent with most previous claims, no evidence was found for evolution in the XLF within $z \leq 0.2 - 0.3$ (Ebeling *et al.* 1998).

We have examined a statistically complete sample of 294 Abell rich clusters within $z \leq 0.09$ using the *ROSAT* All-Sky-Survey (RASS) over the energy band 0.5 – 2 keV as part of a multiwavelength study of nearby galaxy clusters. Unlike most other studies, our sample is purely optically-selected within the criteria for inclusion in Abell's catalog. There is some overlap with both the BCS and XBACS sample, with the primary differences that we have used only Abell's northern catalog (Abell 1958), and our X-ray flux-limit is approximately a factor of 8 lower than the BCS sample. Our sample is larger than the BCS while our volume is nearly 30 times smaller. Given our large sample size, we have reduced statistical errors in the local XLF for $L_X \geq 10^{43} h_{50}^{-2} \text{ ergs sec}^{-1}$ by up to a factor of 2 compared to previous work. Combined with the poor cluster XLF of Burns *et al.* (1996) (BLL96), we examine the composite local XLF over more than 3 orders of magnitude in L_X in order to understand the cosmological constraints imposed by the tight power-law shape noted in BLL96.

In section 2 we describe the sample, the derivation of the local XLF, and discuss the limitations imposed by our sample selection. In section 3 we compare our new XLF with previous work. In section 4 we explore the consequences of the shape of the local XLF with regard to Press-Schechter

¹The National Radio Astronomy Observatory is operated by Associated Universities, Inc., under contract with the National Science Foundation.

analytic predictions of the mass-function and possible constraints on Ω_0 and Λ . We list our conclusions in section 5. We adopt $H_0 = 50 h_{50} \text{ km sec}^{-1}$ and $q_0 = 0.5$ when dealing with the observational data.

2. THE SAMPLE AND DERIVATION OF THE XLF

Our cluster sample is derived from Abell's Northern catalog, and includes all Abell clusters in the range $0.016 \leq z \leq 0.09$ with galactic absorption less than 0.1 magnitudes at R-band ($\log N_H \approx 20.73$). See Voges *et al.* (1999) and Ledlow & Owen (1995) for more details on the sample selection. The total sample includes 294 Abell clusters. All clusters have measured redshifts and we include all richness classes in the sample. We calculate a survey area of $14,155 \text{ deg}^2$ or 34% of the sky. Within our observed volume we find the number density of clusters to be constant as a function of richness class and redshift suggesting that our sample is nearly complete and volume-limited within the limits of Abell's selection criteria. These findings are consistent with those of Briel & Henry (1993) and Mazure *et al.* (1996) with regards to the completeness of Abell's catalog over this redshift regime.

The X-ray luminosity function was derived from images produced by the RASS as described in Voges *et al.* (1999). X-ray luminosities were calculated within a metric aperture of $0.75 h_{50}^{-1} \text{ Mpc}$ in diameter over the energy band 0.5-2 keV assuming a thermal spectrum with $T = 5 \text{ keV}$. Corrections for missing flux were made according to the prescription of Briel & Henry (1993) (using $\beta = 2/3$) to produce a total L_X for each cluster over our *ROSAT* band. The primary effect of using a different β would be to shift the total luminosities to higher or lower values (a larger β results in a smaller correction, thus lower total L_X), while not significantly changing the shape or amplitude of the XLF within the error bars.

Voges *et al.* found a total detection rate of 83% for this sample of Abell clusters. For non-detections, we adopt the 3σ upper-limits given in their Table 1. Because of variations in exposure time (and slight variations in galactic absorption) across the sky with the RASS, each cluster has a different flux-limit, or maximum volume to which the cluster could have been detected. We follow the prescription of Avni & Bahcall (1980), and calculate the observed volume separately for each cluster. The volume is evaluated from $z_{min}=0.016$ to the maximum redshift at which the cluster could have been detected with a 3σ confidence. For clusters with only upper-limits to L_X we set z_{max} equal to the redshift of the cluster. The XLF is then found by calculating $dn(L)/dL$ as the sum over all clusters divided by the maximum search volumes of each cluster. Each binned data point is then found by dividing the above sum by the binwidth (ΔL_X). For the entire sample, we find $\langle V/V_{max} \rangle = 0.56 \pm 0.02$. Error bars on the data points were calculated assuming Poisson statistics following the prescription of Rosati *et al.* (1998).

3. THE X-RAY LUMINOSITY FUNCTION

In Figure 1 we show the differential XLF for our low-redshift cluster sample. Also on this plot are the measurements of BLL96 derived from 49 poor clusters and the BCS sample of Ebeling *et al.* (1998). The steady decline in volume-density observed in our rich cluster sample for

$L_X < 10^{43} h_{50}^{-2} \text{ ergs/sec}$ can be understood from the limitations of Abell's optical selection criteria. Because L_X varies considerably for a given optical richness (Voges *et al.* 1999), there are a significant number of optically poor clusters with L_X in the range of Richness Class 0 clusters which are not in our sample. Thus, our sample is truly volume-limited only for clusters above this cutoff in L_X . Note, however, that for $L_X > 10^{43} \text{ ergs/sec}$, our Abell cluster sample and the BCS sample are in excellent agreement. The BCS also extends to higher L_X because of the larger search volume ($z \leq 0.3$). Our XLF shown in Figure 1 is also consistent with that of Edge *et al.* (1990) and Briel & Henry (1993).




Fig. 1. The X-ray luminosity function derived from our low-redshift ($z \leq 0.09$) Abell cluster sample (solid circles), the poor-cluster data points from Burns *et al.* (1996) (open squares), and the XLF from the Brightest-Cluster Sample (BCS) of Ebeling *et al.* (1998) (open circles).

The local, differential XLF is remarkably well represented by a power-law over more than three orders of magnitude in L_X . The high luminosity break in the XLF occurs at $> 10^{45} h_{50}^{-2} \text{ ergs/sec}$, and can be seen when we include the highest luminosity point from the BCS sample. Using the combined XLF of BLL96 and our new determination of the local rich-cluster XLF (for $L_X > 10^{43} h_{50}^{-2} \text{ ergs/sec}$), we find a power-law fit of the form $\phi(L) = K L_{44}^{-\alpha}$ where L_{44} is the X-ray luminosity in units of $10^{44} \text{ ergs sec}^{-1}$ and K is in units of $10^{-7} \text{ Mpc}^{-3} L_{44}^{\alpha-1}$. We find best-fit values of $\alpha = 1.83 \pm 0.04$ and $K = 2.35 \pm 0.24$. For completeness, we also fit a Schechter function after including the highest- L_X point from the BCS. For a fit of the form: $\frac{dN}{dL_X} = A \exp(-L_X/L_X^*) L_X^{-\alpha}$, we find $A = (2.93 \pm 0.14) \times 10^{-7} (\text{Mpc}^{-3} L_{44}^{\alpha-1})$, $L_{X(0.5-2\text{keV})}^* = 5.49 \pm 0.39 (10^{44} \text{ ergs/sec})$, and $\alpha = 1.77 \pm 0.01$. These values are consistent within the errors to the BCS, the RDCS XLF (Rosati *et al.* 1998) out to $z=0.6$, and the Southern SHARC survey (Burke *et al.* 1997) for $0.3 < z < 0.7$. Note that these results do not conflict with the claimed negative evolution in the XLF observed by Henry *et al.* (1992), and most recently by Vikhlinin *et al.* (1998) at the highest luminosities.

As noted by BLL96, the remarkable power-law shape over such a large range in L_X suggests a continuity in that the bulk X-ray properties of poor clusters must not be fundamentally different from richer systems. We explore the

consequences of this result in the next section.

4. DERIVATION OF THE THEORETICAL XLF

In order to assess the constraints our local XLF imposes on cosmological models, we compare it with various analytic predictions. We proceed by using the Press-Schechter (PS) formalism (e.g., Press & Schechter 1974; Bond *et al.* 1991) to construct theoretical mass functions and then convert these to XLFs assuming a form for the X-ray mass-to-light ratio (c.f., Evrard & Henry 1991; hereafter EH91).

We begin with the set of cosmological models whose parameters are listed in Table 1. These models form a representative sample of current views as they include open and flat universes spanning a range in Ω_o . For each model, the rms density fluctuation on $8h^{-1}\text{Mpc}$ scales (σ_8) was determined from the $\sigma_8 - \Omega_o$ relation of Viana & Liddle (1996) which, in turn, was fixed by the local number density of 7 keV clusters. The Hubble constant was chosen to give an age for the Universe of roughly 12.5 Gyrs (consistent with globular cluster age determinations; e.g., Chaboyer *et al.* 1998). For each model we list the relative contributions of matter (Ω_o), baryonic matter (Ω_b), and the cosmological constant (Ω_Λ) to the overall energy density. Power spectra for all the models were generated using the code described in Klypin & Holtzman (1997) and then PS mass functions (with $\delta_c = 1.3$) were computed at $z = 0$.

Our PS mass functions can be converted to XLFs by assuming a form for the mass-luminosity relation and correcting to our bandpass. We assume the bolometric X-ray luminosity is related to cluster mass as $L_{bol} = cM^p$ and will later fit for the parameters c and p . There exist at least two theoretical predictions for the value of the exponent p . The self-similar model of Kaiser (1986), derived assuming a power-law initial perturbation spectrum and purely adiabatic gas physics, predicts $p = 4/3$ but it is well known that this fails to give the correct shape for the XLF (e.g., EH91; see also below). However, pre-heating of the ICM at an early epoch (possibly by galaxy formation) results in a different scaling relation and also resolves several discrepancies between theoretical and observational results concerning evolution in the XLF (e.g., Evrard 1990, Navarro, Frenk & White 1995). For the case of a constant entropy core, EH91 derived a scaling which implies $p = (10\beta - 3)/3\beta$ where $\beta = \mu m_p \sigma^2 / kT$ is the usual ratio of dark matter to gas ‘temperatures’.

We correct our bolometric luminosities to the 0.5-2 keV bandpass by calculating temperatures and applying a correction appropriate for a plasma with a metallicity of $Z=0.3Z_\odot$. Specifically, the temperature corresponding to a given mass can be calculated from the analytic M-T relation derived from the virial theorem (e.g., Bryan & Norman 1998): $kT = \frac{1.39}{\beta} \left(\frac{M}{10^{15} M_\odot} \right)^{2/3} \Delta_c^{1/3} h^{2/3} \text{ keV}$ where Δ_c is the current density contrast within the cluster virial radius. The luminosity in our bandpass is then calculated by applying the usual bremsstrahlung correction factor as well as a multiplicative factor to account for the presence of metals (Bryan & Norman 1998; eqn. 21).

Using the relation for L_{bol} and the bandpass correction, we converted our PS mass functions to differential luminosity functions and made χ -squared fits to a subset of the observational data. The observational points used in the fits are all four poor cluster points (BLL96), the five

highest luminosity Abell cluster points and the highest luminosity BCS point from Figure 1. We first set $\beta = 1$ in the $M - T$ relation and fit for c and p . The fitted value for p is included in Table 1 and examples of two of the fits are shown in Figure 2. The dashed curve in Figure 2a is the best fit when the exponent is kept fixed at the analytic prediction $p = 4/3$. Clearly, the shape of the XLF derived using this prediction is in gross disagreement with the observed function. Figure 2b also shows the importance of the low and/or high-luminosity data points. If only our five Abell cluster data points are used (dotted line), the fitted value of p increases by at least 0.2 in all cases (from $p = 3.18$ to $p = 3.88$ in this case). We get virtually identical results if we redo our fits without the BCS point whereas dropping the poor cluster points results in slightly greater discrepancies.




Fig. 2. Fits to the observed XLF for two models. The solid lines are the best fits to a subset of the observational XLF points for models ACDM1 (left panel) and ACDM5 (right panel). The dashed line in the left panel is the best-fit to the data assuming $p=4/3$ as predicted by the self-similar analytic models. The dotted line in the right panel represents the best-fit to the data when the BCS and poor cluster points are ignored.

If we invoke the constant entropy core model of EH91, then the exponent in the mass-luminosity relation is actually a function of β ($p = (10\beta - 3)/3\beta$). In this case, we fit for c and β and find the values listed in Table 1. Interestingly, only the models with $0.1 < \Omega_o < 0.4$ are consistent with the expected value $\beta \sim 1$. [A recent observational analysis found $\beta = 0.94 \pm 0.08$ (Lubin & Bahcall 1993) which is in good agreement with numerical results (e.g., Eke, Navarro, & Frenk 1998).] Thus, if the constant entropy core model of EH91 applies, the present-day XLF observations suggest a low-density universe but cannot distinguish between open and flat cases.

5. CONCLUSIONS

Starting from an optically-selected, statistical sample of Abell clusters, we have made a new determination of the local XLF to compare to previous work and more distant cluster samples. Our cluster sample is larger than all previous studies, and is contained within a smaller volume. For this reason, we have reduced statistical uncertainties in the local XLF by nearly a factor of two for a limited range in L_X ($L_X > 10^{43} h_{50}^{-2} \text{ ergs/sec}$). It is only for $L_X < 10^{43} \text{ ergs/sec}$ that incompleteness due to the opti-

TABLE 1
COSMOLOGICAL MODEL PARAMETERS

Model	Age ^a	H_0^b	Ω_o	Ω_b	Ω_Λ	σ_8	p	β
OCDM1	12.5	70	0.1	0.026	0.0	1.467	$1.76^{+0.17}_{-0.14}$	$0.66^{+0.07}_{-0.05}$
OCDM2	12.7	65	0.2	0.030	0.0	1.162	$2.20^{+0.17}_{-0.17}$	$0.88^{+0.10}_{-0.10}$
OCDM3	12.2	65	0.3	0.030	0.0	1.004	$2.50^{+0.24}_{-0.20}$	$1.14^{+0.41}_{-0.18}$
OCDM4	12.7	60	0.4	0.035	0.0	0.897	$2.69^{+0.25}_{-0.32}$	$1.47^{+0.64}_{-0.33}$
OCDM5	12.3	60	0.5	0.035	0.0	0.817	$2.86^{+0.21}_{-0.21}$	$1.90^{+0.5}_{-0.5}$
OCDM6	11.6	60	0.7	0.035	0.0	0.702	$3.22^{+0.29}_{-0.29}$	$4.0^{+2.0}_{-2.1}$
Λ CDM1	13.1	80	0.2	0.020	0.8	1.478	$2.14^{+0.18}_{-0.15}$	$0.84^{+0.08}_{-0.08}$
Λ CDM2	11.8	80	0.3	0.020	0.7	1.160	$2.49^{+0.21}_{-0.17}$	$1.17^{+0.33}_{-0.19}$
Λ CDM3	12.4	70	0.4	0.026	0.6	0.901	$2.77^{+0.27}_{-0.22}$	$1.62^{+0.63}_{-0.37}$
Λ CDM4	12.5	65	0.5	0.030	0.5	0.864	$2.86^{+0.31}_{-0.21}$	$1.90^{+0.75}_{-0.50}$
Λ CDM5	12.2	60	0.7	0.035	0.3	0.719	$3.18^{+0.31}_{-0.27}$	$3.5^{+2.4}_{-1.6}$

^a current age of universe in Gyrs

^b Hubble constant in units of km/s/Mpc

cal selection of our sample is apparent. The observed incompleteness is not a failing in Abell's catalog, but rather results from the contribution of poor clusters and groups below Abell's richness limit.

Combined with the poor-cluster XLF of BLL96, we have examined the local XLF over nearly three orders of magnitude in L_X . We find that the local XLF is remarkably well represented by a power-law over nearly this entire range in L_X . This is significant evidence that hierarchical formation results in similar cluster properties over a large range in L_X and mass. Including the brightest L_X clusters from the BCS sample which fall above the break in the XLF at $L_X > 10^{45} h_{50}^{-2} \text{ ergs/sec}$, we also performed a Schechter-function fit which is in good agreement with other recent surveys to much higher redshift ($z < 0.7$), confirming a lack of significant evolution at these luminosities.

We have used our new *local* XLF to derive a constraint on Ω_0 . This would appear to contradict a common claim that the $\sigma_8 - \Omega_0$ degeneracy can be broken only by including the evolution with redshift (*e.g.* Bahcall & Fan 1998). In fact, PS mass functions for combinations of σ_8 and Ω_0 that satisfy a $\sigma_8 - \Omega_0$ constraint differ in shape. Borgani *et al.* (1999) have recently used the shape of the local XLF in

order to constrain $\sigma_8 - \Omega_0$ and the shape of the L-T relation. Including clusters at higher redshift, they concluded that $\Omega_0 = 0.4^{+0.3}_{-0.2}$ for open models, and $\Omega_0 \leq 0.6$ for flat models assuming no evolution in the L-T relation; both of which are consistent with our results. In this work, we have used the shape of the local XLF, the local number density of 7 keV clusters, and the PS formalism in order to constrain the cluster M-L relation; $L_X \propto M^p$. There is a clear trend for p to increase with Ω_0 (see also Mathiesen & Evrard 1998). None of the theoretical models are consistent with the analytic prediction $p = 4/3$ from Kaiser (1986). If we adopt the constant core-entropy model of EH91, and the additional constraint that $\beta \approx 1$, the shape of the local XLF suggests that $0.1 \leq \Omega_0 \leq 0.4$, with no constraint on Λ .

Acknowledgements

This work was supported in part by NASA grants NAG5-6739 and NAGW-3152, and NSF grant AST-9896039. We thank Anatoly Klypin and Jon Holtzman for use of their code and useful discussions. We also thank Neta Bahcall and an anonymous referee for helpful suggestions. JOB and FNO thank MPE for their hospitality during several visits.

REFERENCES

- Abell, G.O. 1958, ApJS, 31, 211
Avni, Y. & Bahcall, J.N., 1980, ApJ, 235, 694
Bahcall, N.A. & Fan, X. 1998, ApJ, 504, 1
Bond, J.R., Cole, S., Efstathiou, G., & Kaiser, N. 1991, ApJ, 379, 440
Borgani, S., Rosati, P., Tozzi, P., & Norman, C. 1999, astro-ph/9901017
Briel, U.G. & Henry, J.P. 1993, A&A, 278, 379
Bryan, G. & Norman, M. L. 1998, ApJ, 495, 80
Burke, D.J., Collins, C.A., Sharples, R.M., Romer, A.K., Holden, B.P., & Nichol, R.C., 1997, ApJ, 488, L83
Burns, J.O., Ledlow, M.J., Loken, C., Klypin, A., Voges, W., Bryan, G.L., Norman, M.L., & White, R.A. 1996, ApJ, 467, L49 (BLL96)
Chaboyer, B., Demarque, P., Kernan, P.J., & Krauss, L.M. 1998, ApJ, 494, 96
Ebeling, H., Edge, A.C., Böhringer, H., Allen, S.W., Crawford, C.S., Fabian, A.C., Voges, W., & Huchra, J.P. 1998, MNRAS, 301, 881
Ebeling, H., Edge, A.C., Fabian, A.C., Allen, S.W., Crawford, C.S., & Böhringer, H. 1997, ApJ, 479, L101
Ebeling, H., Voges, W., Böhringer, H., Edge, A.C., Huchra, J.P., & Briel, U.G. 1996, MNRAS, 283, 1103
Ebeling, H., Voges, W., Böhringer, H., & Edge, A.C. 1993, A&A, 275, 360
Edge, A.C., Stewart, G.C., Fabian, A.C., & Arnaud, K.A. 1990, MNRAS, 245, 559
Eke, V.R., Navarro, J.F., & Frenk, C.S. 1998, ApJ, 503, 569
Evrard, A.E. & Henry, J.P. 1991, ApJ, 383, 95 (EH91)
Evrard, A.E. 1990, ApJ, 363, 349
Henry, J.P., Gioia, I.M., Maccacaro, T., Morris, S.L., Stocke, J.T., & Wolter, A. 1992, ApJ, 386, 408
Jones, L.R., Scharf, C., Ebeling, H., Perlman, E., Wegner, G., Malkan, M., & Horner, D. 1998, ApJ, 495, 100
Kaiser, N. 1986, MNRAS, 219, 785
Klypin, A. & Holtzman, J. 1997, astro-ph/9712217
Ledlow, M.J. & Owen, F.N. 1995, AJ, 109, 853
Lubin, L.M. & Bahcall, N.A. 1993, ApJ, 415, L17
Mathiesen, B. & Evrard, A.E. 1998, MNRAS, 295, 769
Mazure, A., Katgert, P., den Hartog, R., Biviano, A., Dubath, P., Escalera, E., Focardi, P., Gerbal, D., Giuricin, G., Jones, B., Le Fevre, O., Moles, M., Perea, J., & Rhee, G. 1996, A&A, 310, 31
Navarro, J.F., Frenk, C.S., & White, S.D.M. 1995, MNRAS, 275, 720
Press, W.H. & Schechter, P. 1974, ApJ, 187, 425
Rosati, P., Ceca, R.D., Norman, C., & Giacconi, R. 1998, ApJ, 492, L21
Viana, P.T.P. & Liddle, A.R. 1996, MNRAS, 281, 323
Vikhlinin, A., McNamara, B.R., Forman, W., Jones, C., Quintana, Q., & Hornstrup, A. 1998, astro-ph/9803101
Voges, W., Ledlow, M.J., Burns, J.O., Owen, F.N., & Smith, C. 1999, submitted to AJ